

Renormalization group parameter evolution of the minimal supersymmetric standard model with R -parity violation

Marek Gózdź* and Wiesław A. Kamiński†

Theoretical Physics Department, Maria Curie-Skłodowska University, Lublin, Poland

A comparison of spectra obtained using the 1-loop MSSM and 2-loop R_p -MSSM renormalization group equations is presented. Influence of higher loop corrections and R -parity violating terms is discussed. Some numerical constraints on the R -parity violating parameters are also given.

PACS numbers: 11.10.Hi, 11.30.Pb, 12.60.Jv, 23.40.Bw

Keywords: R -parity violation, MSSM, supersymmetry, renormalization group, neutrinoless double beta decay

I. INTRODUCTION

The recent observations of neutrino oscillations [1, 2, 3] and reported, although not confirmed, discovery of neutrinoless double beta decay [4] give strong motivation for studying some aspects of physics beyond the Standard Model (SM). One of the most popular approach to such a physics is the incorporation of supersymmetry, which leads, in the simplest case, to the Minimal Supersymmetric Standard Model (MSSM) [5]. This framework is very attractive for theorists since it is the simplest supersymmetric model possible that leads to unification of the coupling constants. It also provides a natural way to incorporate of lepton and baryon number violating processes, which is impossible in SM.

It was widely approved to neglect the terms violating lepton and baryon numbers in actual calculations, due to smallness of the relevant coupling constants [6, 7, 8]. However, even if perfectly valid in low energies, this assumption could have lead to erroneous predictions in high energy limits. We show by explicit simulations that this approach is justified in most cases.

The paper is organized as follows. In the next section we present the model. We also explain our conventions and notations here. In Section III the results are presented and discussed. A short conclusion is given at the end.

II. THEORY

The minimal (in the particle and interaction content) supersymmetric extension of SM means introducing a superpartner for each particle and adding a second Higgs doublet (together with its higgsinos).

We closely follow the conventions used in [9, 10]. The MSSM superpotential can be written in the form

$$W^{MSSM} = \epsilon_{ab}[(\mathbf{Y}_E)_{ij}L_i^a H_1^b \bar{E}_j + (\mathbf{Y}_D)_{ij}Q_i^{ax} H_1^b \bar{D}_{jx} + (\mathbf{Y}_U)_{ij}Q_i^{ax} H_2^b \bar{U}_{jx} + \mu H_1^a H_2^b], \quad (1)$$

where $\mathbf{Y}$'s are 3×3 Yukawa matrices. L and Q are the $SU(2)$ left-handed doublets while $\bar{E}$, $\bar{U}$ and $\bar{D}$ denote the right-handed lepton, up-quark and down-quark $SU(2)$ singlets, respectively. H_1 and H_2 mean two Higgs doublets. We have introduced color indices $x, y, z = 1, 2, 3$, generation indices $i, j, k = 1, 2, 3$ and the $SU(2)$ spinor indices $a, b, c = 1, 2$.

Since supersymmetry is not observed, we need to introduce some mechanism to break it. The simplest possibility is the so-called soft breaking given by the Lagrangian

$$\mathcal{L}^{soft} = \epsilon_{ab}[(\mathbf{A}_E)_{ij}l_i^a h_1^b \bar{e}_j + (\mathbf{A}_D)_{ij}q_i^{ax} h_1^b \bar{d}_{jx} + (\mathbf{A}_U)_{ij}q_i^{ax} h_2^b \bar{u}_{jx} + B\mu h_1^a h_2^b], \quad (2)$$

where lowercase letters stand for scalar components of respective chiral superfields, and 3×3 matrices $\mathbf{A}$ as well as $B\mu$ are the soft breaking coupling constants. We introduce also a scalar mass term of the form

$$\mathcal{L}^{mass} = \mathbf{m}_{H_1}^2 h_1^\dagger h_1 + \mathbf{m}_{H_2}^2 h_2^\dagger h_2 + q^\dagger \mathbf{m}_Q^2 q + l^\dagger \mathbf{m}_L^2 l + u \mathbf{m}_U^2 u^\dagger + d \mathbf{m}_D^2 d^\dagger + e \mathbf{m}_E^2 e^\dagger. \quad (3)$$

Looking at (1) it is clearly visible, that W^{MSSM} preserves a discrete symmetry called R -parity and defined as

$$R_p = (-1)^{3B+L+2S},$$

where B , L , and S are the baryon, lepton and spin numbers, respectively. From the definition follows that ordinary particles have $R_p = +1$ whereas supersymmetric (SUSY) particles have $R_p = -1$. The consequence of R_p conservation is a stable lightest SUSY particle, which can be a serious candidate for cold dark matter, and a stable proton.

However, nothing motivates theoretically R_p conservation, especially that in order to avoid rapid proton decay, at least the lepton number parity $L_p = (-1)^{L+2S}$ or the baryon number parity $B_p = (-1)^{3B+2S}$ must be conserved, *but not necessarily both*. It follows that we can add new terms to the superpotential, which violate separately the lepton and baryon number:

$$W^{R_p} = \epsilon_{ab} \left[\frac{1}{2} (\mathbf{A}_{E^k})_{ij} L_i^a L_j^b \bar{E}_k + (\mathbf{A}_{D^k})_{ij} L_i^a Q_j^{xb} \bar{D}_{kx} \right] + \frac{1}{2} \epsilon_{xyz} (\mathbf{A}_{U^i})_{jk} \bar{U}_i^x \bar{D}_j^y \bar{D}_k^z + \epsilon_{ab} \kappa^i L_i^a H_2^b. \quad (4)$$

*Electronic address: mgozdz@kft.umcs.lublin.pl

†Electronic address: kaminski@neuron.umcs.lublin.pl

The full superpotential becomes now

$$W = W^{MSSM} + W^{\mathcal{R}_p} \quad (5)$$

and defines a new model, often called $\mathcal{R}_p$ -MSSM. It introduces nine new Yukawa coupling constants but, as mentioned above, proton stability requires to reject either lepton number or baryon number violating terms. If one wants to apply this model to the description of the neutrinoless double beta decay

$$A(Z, N) \rightarrow A(Z + 2, N - 2) + 2e^-,$$

in which the lepton number is violated by two units, the matrices $\mathbf{A}_{U^i}$ have to be set equal to zero. In this paper we will not do that in order to stay completely general and to be able to count contributions coming from all sources.

All the parameters (coupling constants and masses) of $\mathcal{R}_p$ -MSSM are running parameters in the sense of renormalization group, that is they obey a set of equations of the form

$$\mu \frac{dx}{dt} = \frac{1}{16\pi^2} \beta_x^{(1)} + \frac{1}{(16\pi^2)^2} \beta_x^{(2)} + \dots, \quad (6)$$

where $t = \log E$ and E is the renormalization scale; $\beta_x^{(n)}$ is the n -loop renormalization group beta function for the parameter x . Our previous calculations [8, 11, 12] were based on 1-loop MSSM renormalization group equations because we roughly estimated that the corrections coming from R -parity violation are of the order of few percent. We check this assumption in detail in the next section.

III. RESULTS

The relevant 1- and 2-loop equations for MSSM can be found in [9] with $\mathcal{R}_p$ -MSSM corrections in [10]. Our procedure of running the parameters consists of few steps. We take into account mass thresholds where SUSY particles start to contribute [13] and use the SM 2-loop renormalization group equations (RGE) below appropriate threshold and 2-loop $\mathcal{R}_p$ -MSSM RGE above it. Initially all the thresholds are set to 1 TeV and are dynamically modified during the running of mass parameters. After evolving the dimensionless couplings from the electroweak scale m_Z up to $m_{GUT} \sim 10^{16}$ GeV, we unify the masses of gauginos, sfermions and squarks to be $m_0 = 500$ GeV. The trilinear soft couplings at m_{GUT} are set according to formula

$$\mathbf{A}_i = A_0 \mathbf{Y}_i,$$

with $A_0 = 500$. Next, all the quantities are evolved down to m_Z . The numbers used in simulation serve as an example. Our investigation showed, that the same conclusions can be drawn for a wide range of m_0 and A_0 .

TABLE I: Mass matrices of supersymmetric particles (third family) and values of the coupling constants, calculated using 1-loop MSSM and 2-loop $\mathcal{R}_p$ -MSSM renormalization group equations. All supersymmetric masses are unified (500 GeV) at the GUT scale 10^{16} GeV. Other MSSM parameters were $\tan \beta = 5$ and positive sign of μ .

	1-loop GUT MSSM	2-loop GUT $\mathcal{R}_p$ -MSSM	1-loop m_Z MSSM	2-loop m_Z $\mathcal{R}_p$ -MSSM
$\frac{5}{3}\alpha_1$	0.0396	0.0398	0.0169	0.0169
α_2	0.0386	0.0402	0.0338	0.0338
α_3	0.0387	0.0396	0.1177	0.1171
Y_τ	0.0356	0.0371	0.0527	0.0527
Y_b	0.0337	0.0343	0.0937	0.0937
Y_t	0.6356	0.6388	0.9598	0.9598
A_τ/Y_τ	500	500	168	182
A_b/Y_b	500	500	-1131	-1170
A_t/Y_t	500	500	-853	-861
μ	500	500	761	805
$B\mu$	500	500	-247	-248
M_1	500 GeV	500 GeV	210 GeV	210 GeV
M_2	500 GeV	500 GeV	415 GeV	414 GeV
M_3	500 GeV	500 GeV	1290 GeV	1308 GeV
m_{H_1}	500 GeV	500 GeV	604 GeV	598 GeV
$\pm m_{H_2} $	500 GeV	500 GeV	-738 GeV	-782 GeV
m_{udL}^2	500 GeV	500 GeV	1282 GeV	1284 GeV
m_{csL}^2	500 GeV	500 GeV	1281 GeV	1284 GeV
m_{tbL}^2	500 GeV	500 GeV	1155 GeV	1163 GeV
m_{uR}^2	500 GeV	500 GeV	1242 GeV	1252 GeV
m_{cR}^2	500 GeV	500 GeV	1242 GeV	1252 GeV
m_{tR}^2	500 GeV	500 GeV	963 GeV	984 GeV
m_{dR}^2	500 GeV	500 GeV	1237 GeV	1249 GeV
m_{sR}^2	500 GeV	500 GeV	1237 GeV	1249 GeV
m_{bR}^2	500 GeV	500 GeV	1234 GeV	1247 GeV
$m_{\nu eL}^2$	500 GeV	500 GeV	612 GeV	607 GeV
$m_{\nu \mu L}^2$	500 GeV	500 GeV	612 GeV	607 GeV
$m_{\nu \tau L}^2$	500 GeV	500 GeV	612 GeV	606 GeV
m_{eR}^2	500 GeV	500 GeV	536 GeV	532 GeV
$m_{\mu R}^2$	500 GeV	500 GeV	536 GeV	532 GeV
$m_{\tau R}^2$	500 GeV	500 GeV	534 GeV	531 GeV

The values of Yukawa couplings $\mathbf{Y}$ at m_Z are given by lepton and quark mass matrices M [5]

$$\begin{aligned} M_U &= \langle H_2^0 \rangle \mathbf{S}_{U_R} \mathbf{Y}_U^T \mathbf{S}_{U_L}^\dagger, \\ M_D &= \langle H_1^0 \rangle \mathbf{S}_{D_R} \mathbf{Y}_D^T \mathbf{S}_{D_L}^\dagger, \\ M_E &= \langle H_1^0 \rangle \mathbf{S}_{E_R} \mathbf{Y}_E^T \mathbf{S}_{E_L}^\dagger, \end{aligned} \quad (7)$$

where $\langle H_1^0 \rangle$ and $\langle H_2^0 \rangle$ are the neutral Higgs vacuum expectation values. $\mathbf{S}$ matrices perform diagonalization so that one obtains eigenstates in the mass representation. However, we had to choose some low energy values for the remaining R -parity violating couplings $\mathbf{A}$. There is no hint concerning their actual numerical values except that they should be small, since lepton number violating processes are very rare. We checked several possibilities and found, that in order to obtain physically reliable re-




FIG. 1: R -parity violating Yukawa couplings evolution. All were set to the same value at the m_Z scale and evaluated using the 2-loop $\mathcal{R}_p$ -MSSM RGE to the m_{GUT} scale.

sults, all these couplings should obey

$$(\Lambda)_{ii} \leq 0.015, \quad (i = 1, 2, 3), \quad (8)$$

otherwise high vacuum expectation values are generated from the down quark mass evolution. What is more, values of other parameters nearly did not change for various Λ 's. From [12] we can make use of the following relation

$$(\Lambda_{D^1})_{11} \leq \xi \left(\frac{10^{24} y}{T_{1/2}^{0\nu 2\beta - exp}} \right)^{-1/4}. \quad (9)$$

Using the ξ parameter for ^{76}Ge isotope for the unification mass 500 GeV, and inserting the experimentally known half-life given by the Heidelberg–Moscow collaboration [4] $T_{1/2}^{0\nu 2\beta(HMexp)} = (0.8 - 18.3) \times 10^{25} y$, we obtain

$$(\Lambda_{D^1})_{11} \leq 0.275, \quad (10)$$

which is much weaker than our constraint (8). If one uses limit (8) the half-life to be of the order of $\sim 1.3 \times 10^{30} y$ which is rather an overestimation. On the other hand, the newest data from the IGEX Collaboration [14] sets the lower bound on the half-life in germanium to be $T_{1/2}^{0\nu 2\beta(IGEXexp)} > 1.57 \times 10^{25} y$ without bounds from above.

Values for various parameters in both 1-loop MSSM and 2-loop $\mathcal{R}_p$ -MSSM cases are given in Tab.I. The $SU(3)_c \times SU(2)_L \times U(1)_Y$ dimensionless coupling constants g_i are represented here in the usual way by $\alpha_i = g_i^2/4\pi$. Whenever the parameter has a matrix form, it has been diagonalized and the entry $\{33\}$ is listed in the table. Masses are given respectively for gauginos, left-handed Higgs doublets, left-handed squark doublets, right-handed up- and down-type squark singlets, left-handed slepton doublets, and right-handed slepton singlets, with respect to the $SU(2)$ group.

In general, as expected, all parameters are changed by at most few percent. One should keep in mind that the new results include two corrections: the 2-loop terms and the terms connected with R -parity violation. Our investigation shows, that the differences are mainly driven by the 2-loop corrections, whereas the R -parity terms change the results on the third decimal place.

This situation can be understood from the evolution of R_p -violating couplings. In Fig.1 an example of running of Λ coupling constants is shown. It is essential, that their values decrease with increasing energy. As a consequence the contributions coming from Λ 's can be safely neglected in actual calculations. It is, however, a surprising result, since normally one would expect to see exotic processes at high energies. Fig.1 clearly shows, that to observe a lepton number violating process, like the neutrinoless double beta decay, one should search for

it in low temperatures rather than in accelerators.

IV. FINAL REMARKS

In conclusion, we have presented the spectrum of supersymmetric particles and coupling constants using the 1-loop MSSM and 2-loop $\overline{R}_p$ -MSSM renormalization group equations. Inclusion of the R -parity violating terms does not change the results significantly. On the other hand we confirmed the well known rule that the 2-loop corrections should be used in all cases, where quantitative rather than qualitative features are important.

Acknowledgments

This research was partially supported through a European Community Marie Curie Fellowship. One of us

(MG) would like to thank the staff of Marie Curie Trainig Site, located in the European Center for Theoretical Studies in Nuclear Physics and Related Areas (ECT*) in Trento, Italy, for kind hospitality during Summer 2002.

Disclaimer: The authors are solely responsible for the information communicated, published or disseminated. It does not represent the opinion of the Community. The Community is not responsible for any use that might be made of data appearing therein.

-
- [1] Q. R. Ahmed *et al.* (the SNO Collaboration), Phys. Rev. Lett. **89**, 011301 (2002).
 - [2] S. H. Ahn *et al.*, Phys. Rev. Lett. **90**, 041801 (2003).
 - [3] J. Detwiler, G. G., N. Tolich, and Y. Uchida, Phys. Rev. Lett. **89**, 191802 (2002).
 - [4] H. V. Klapdor-Kleingrothaus, A. Dietz, H. L. Harney, and I. Krivosheina, Mod. Phys. Lett. A **16**, 2409 (2001).
 - [5] P. Cho, M. Misiak, and D. Wyler, Phys. Rev. D **54**, 3329 (1996).
 - [6] M. Hirsh, H. V. Klapdor-Kleingrothaus, and S. G. Kovalenko, Phys. Rev. D **53**, 1329 (1996).
 - [7] O. Haug, J. D. Vergados, A. Faessler, and S. G. Kovalenko, Phys. Atom. Nucl. **63**, 1078 (2000).
 - [8] A. Wodecki and W. A. Kamiński, Phys. Rev. C **59**, R1232 (1999).
 - [9] S. P. Martin and M. T. Vaughn, Phys. Rev. D **50**, 2282 (1994).
 - [10] B. C. Allanach, A. Dedes, and H. K. Dreiner, Phys. Rev. D **60**, 056002 (1999).
 - [11] A. Wodecki, W. A. Kamiński, and S. Pagerka, Phys. Lett. B **413**, 342 (1997).
 - [12] A. Wodecki, W. A. Kamiński, and F. Šimkovic, Phys. Rev. D **60**, 11507 (1999).
 - [13] G. L. Kane, C. Kolda, L. Roszkowski, and J. D. Wells, Phys. Rev. D **49**, 6173 (1994).
 - [14] C. E. Aalseth *et al.*, Phys. Rev. D **65**, 092007 (2002).